

THREE POSITIVE SOLUTIONS FOR A CLASS OF ELLIPTIC SYSTEMS IN ANNULAR DOMAINS

JOÃO MARCOS DO Ó¹, SEBASTIÁN LORCA² AND PEDRO UBILLA³

¹*Departamento de Matemática, Universidade Federal da Paraíba,
58059-900, João Pessoa, PB, Brazil (jmbo@mat.ufpb.br)*

²*Departamento de Matemática, Universidad de Tarapacá,
Casilla 7-D, Arica, Chile (slorca@uta.cl)*

³*Universidad de Santiago de Chile, Casilla 307,
Correo 2, Santiago, Chile (pubilla@usach.cl)*

(Received 12 November 2003)

Abstract In this paper we study the existence and multiplicity of positive radial solutions for a class of semilinear elliptic systems in bounded annular domains with non-homogeneous boundary conditions by the use of a fixed-point theorem of cone expansion/compression type.

Keywords: elliptic systems; annular domains; positive radial solutions; multiplicity; fixed points

2000 *Mathematics subject classification:* Primary 35J65; 34B15; 34B18

1. Introduction

In this paper we study the existence and multiplicity of positive radial solutions for a class of semilinear elliptic systems in bounded annular domains. In fact, given real numbers $0 < r_1 < r_2$, we consider the system

$$\left. \begin{aligned} -\Delta u &= a_2 h(|x|, u, v) && \text{in } r_1 < |x| < r_2, \\ -\Delta v &= b_2 k(|x|, u, v) && \text{in } r_1 < |x| < r_2, \\ (u, v) &= (0, 0) && \text{on } |x| = r_1, \\ (u, v) &= (a_1, b_1) && \text{on } |x| = r_2, \end{aligned} \right\} \quad (\text{S})$$

where $(a_1, b_1) \in [0, +\infty)^2 \setminus \{(0, 0)\}$, $(a_2, b_2) \in (0, +\infty)^2$ and the nonlinearities h and k satisfy the following hypotheses.

(H₀) $h, k : [r_1, r_2] \times [0, +\infty)^2 \rightarrow [0, +\infty)$ are continuous functions such that there exist continuous functions, which are non-decreasing in the last two variables h_1 and k_1 , satisfying

$$\left. \begin{aligned} h_1(r, u, v) &\leq h(r, u, v) \leq c_h h_1(r, u, v), \\ k_1(r, u, v) &\leq k(r, u, v) \leq c_k k_1(r, u, v), \end{aligned} \right\} \quad (1.1)$$

where c_h and c_k are positive constants. We recall that ℓ is a non-decreasing function in the last two variables if $\ell(r, u_1, v_1) \leq \ell(r, u_2, v_2)$ whenever $u_1 \leq u_2$ and $v_2 \leq v_1$.

(H₁) There exist $\sigma_1, \sigma_2, \omega_1, \omega_2 \in (r_1, r_2)$ with $\sigma_1 < \sigma_2$ and $\omega_1 < \omega_2$ such that

$$h(r, u, v) > 0 \quad \text{for each } (r, u, v) \in [\sigma_1, \sigma_2] \times ([0, +\infty)^2 \setminus \{(0, 0)\}) \tag{1.2}$$

or

$$k(r, u, v) > 0 \quad \text{for each } (r, u, v) \in [\omega_1, \omega_2] \times ([0, +\infty)^2 \setminus \{(0, 0)\}). \tag{1.3}$$

Let us introduce the following notation:

$$\ell_0 := \lim_{|(u,v)| \rightarrow 0} \frac{\ell(r, u, v)}{|(u, v)|} \quad \text{and} \quad \ell_\infty := \lim_{|(u,v)| \rightarrow +\infty} \frac{\ell(r, u, v)}{|(u, v)|}.$$

(H₂) $h_0 = k_0 = 0$, uniformly in $[r_1, r_2]$ (superlinear at origin).

(H₃) $h_\infty = k_\infty = 0$, uniformly in $[r_1, r_2]$ (sublinear at infinity).

We now state our main result.

Theorem 1.1. *Assume that $h(r, u, v)$ and $k(r, u, v)$ satisfy (H₀)–(H₃). Then the following hold.*

- (i) System (S) has at least one positive solution for all $(a_1, b_1) \in [0, +\infty)^2 \setminus \{(0, 0)\}$ and $(a_2, b_2) \in (0, +\infty)^2$.
- (ii) There exists a positive constant $A > 0$ such that, for all $(a_2, b_2) \in (0, +\infty)^2$ with $\min\{a_2, b_2\} > A$, there exists $\delta > 0$ such that (S) has at least three positive solutions for all $(a_1, b_1) \in [0, +\infty)^2 \setminus \{(0, 0)\}$ with $0 < |(a_1, b_1)| < \delta$.

Our approach to prove Theorem 1.1 relies on fixed-point index theory. The following is a typical example of where our result may be applied:

$$\left. \begin{aligned} -\Delta u &= a_2 c_1(|x|) \hat{h}(u, v) && \text{in } r_1 < |x| < r_2, \\ -\Delta v &= b_2 c_2(|x|) \hat{k}(u, v) && \text{in } r_1 < |x| < r_2, \\ (u, v) &= (0, 0) && \text{on } |x| = r_1, \\ (u, v) &= (a_1, b_1) && \text{on } |x| = r_2, \end{aligned} \right\} \tag{1.4}$$

where \hat{h} and \hat{k} are continuous functions verifying $\hat{h}_0 = \hat{k}_0 = \hat{h}_\infty = \hat{k}_\infty = 0$ and c_1, c_2 are non-negative and non-trivial functions. Moreover, we suppose the following two assumptions.

- (i) $\hat{h}, \hat{k} : [0, +\infty)^2 \rightarrow [0, +\infty)$ are continuous functions such that there exist non-decreasing continuous functions h_1 and k_1 such that

$$\left. \begin{aligned} h_1(u, v) &\leq \hat{h}(u, v) \leq \hat{c}_h h_1(u, v), \\ k_1(u, v) &\leq \hat{k}(u, v) \leq \hat{c}_k k_1(u, v), \end{aligned} \right\} \tag{1.5}$$

where \hat{c}_h and \hat{c}_k are positive constants.

- (ii) $\hat{h}(u, v) > 0$ and $\hat{k}(u, v) > 0$ for each $(u, v) \in ([0, +\infty)^2 \setminus \{(0, 0)\})$.

Thus, according to the assumptions above on functions \hat{h} and \hat{k} , it is not difficult to verify hypotheses (H₀)–(H₃). Therefore, the conclusions of Theorem 1.1 are true.

The study of (1.4) is, in part, motivated by several recent results for elliptic boundary-value problems on annular domains. Among others, we mention [1, 4–6, 8–10] and the references therein. We refer to [5, 6] for homogeneous Dirichlet boundary conditions and [4], as well as [10], for non-homogeneous Dirichlet boundary conditions. In those articles is proved the existence of at least one or two positive solutions, while in the present paper we obtain multiplicity results that include the existence of at least three positive solutions. We would like to mention that we may apply Theorem 1.1 to more general classes of nonlinearities than those considered in (1.4). For example, we may consider in (S) nonlinearities like

$$h(r, u, v) = (a(r)(u^{p_1} + v^{q_1}) + 1)\Phi(b(r)(u^{p_2} + v^{q_2})) \quad \text{and} \quad k(r, u, v) = \frac{(u + v)^{p_3}}{1 + (u + v)^{q_3}},$$

where $p_1, q_1 \in (0, 1)$, $q_2, p_2 \in (1, +\infty)$, $q_3 > p_3 - 1 > 0$, $a, b : [r_1, r_2] \rightarrow [0, +\infty)$ are continuous functions and $\Phi : [0, +\infty) \rightarrow [0, +\infty)$ is a non-decreasing continuous function satisfying

$$\lim_{u \rightarrow 0} \frac{\Phi(u)}{u} = 0 \quad \text{and} \quad \lim_{u \rightarrow +\infty} \frac{\Phi(u)}{u} = \hat{c} > 0.$$

We notice that the functions a, b may vanish in parts of the interval $[r_1, r_2]$. Finally, we observe that hypothesis (H₁) allows us to take nonlinearities h and k without the monotonicity property.

The rest of this paper is organized as follows. Section 2 contains preliminary results and § 3 is devoted to proving our main result, Theorem 1.1.

2. Preliminary results

We give some results which will be necessary to prove Theorem 1.1 in the next section.

The proof of Theorem 1.1 relies on fixed-point index theory in the frame of the ordinary-differential-equation (ODE) technique. Since we are interested in the existence of radial solutions, by applying consecutive changes of variables with $r = |x|$ and

$$t = -\eta_1 r^{2-N} + \eta_2, \quad \text{where} \quad \eta_1 = \frac{(r_1 r_2)^{N-2}}{r_2^{N-2} - r_1^{N-2}} \quad \text{and} \quad \eta_2 = \frac{r_2^{N-2}}{r_2^{N-2} - r_1^{N-2}},$$

we can transform (S) into the following system of second-order ODEs,

$$\left. \begin{aligned} -u'' &= a_2 f(t, u, v, a_1, b_1) && \text{in } (0, 1), \\ -v'' &= b_2 g(t, u, v, a_1, b_1) && \text{in } (0, 1), \\ u(0) &= u(1) = 0, \\ v(0) &= v(1) = 0, \end{aligned} \right\} \tag{R}$$

where here the nonlinearities f and g are given by

$$\left. \begin{aligned} f(t, u, v, a_1, b_1) &= d(t)h\left(\left(\frac{\eta_1}{\eta_2 - t}\right)^{1/(N-2)}, u + ta_1, v + tb_1\right), \\ g(t, u, v, a_1, b_1) &= d(t)k\left(\left(\frac{\eta_1}{\eta_2 - t}\right)^{1/(N-2)}, u + ta_1, v + tb_1\right), \\ d(t) &= (1 - N)^2 \frac{\eta_1^{2/(N-2)}}{(\eta_2 - t)^{2(N-1)/(N-2)}}. \end{aligned} \right\} \quad (2.1)$$

Setting

$$\begin{aligned} f_1(t, u, v, a_1, b_1) &= d(t)h_1\left(\left(\frac{\eta_1}{\eta_2 - t}\right)^{1/(N-2)}, u + ta_1, v + tb_1\right), \\ g_1(t, u, v, a_1, b_1) &= d(t)k_1\left(\left(\frac{\eta_1}{\eta_2 - t}\right)^{1/(N-2)}, u + ta_1, v + tb_1\right), \end{aligned}$$

we observe that, from (1.1) and (2.1),

$$\left. \begin{aligned} f_1(t, u, v, a_1, b_1) &\leq f(t, u, v, a_1, b_1) \leq c_h f_1(t, u, v, a_1, b_1), \\ g_1(t, u, v, a_1, b_1) &\leq k(t, u, v, a_1, b_1) \leq c_k g_1(t, u, v, a_1, b_1). \end{aligned} \right\} \quad (2.2)$$

It is not difficult to show that if the pair (u, v) is a solution of (R), then, for all $t \in [0, 1]$,

$$\left. \begin{aligned} u(t) &= a_2 \int_0^1 K(t, \tau) f(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau, \\ v(t) &= b_2 \int_0^1 K(t, \tau) g(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau, \end{aligned} \right\} \quad (2.3)$$

where $K(t, \tau)$ is Green's function

$$K(t, s) := \begin{cases} t(1 - s) & \text{if } t \leq s, \\ s(1 - t) & \text{if } t > s. \end{cases} \quad (2.4)$$

Let

$$\begin{aligned} A(u, v)(t) &:= a_2 \int_0^1 K(t, \tau) f(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau, \\ B(u, v)(t) &:= b_2 \int_0^1 K(t, \tau) g(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau, \\ F(u, v) &:= (A(u, v), B(u, v)). \end{aligned}$$

Therefore, system (2.3) is equivalent to the fixed-point equation

$$F(u, v) = (u, v)$$

in the usual Banach space

$$X = C([0, 1], \mathbb{R}) \times C([0, 1], \mathbb{R})$$

endowed with the norm $\|(u, v)\| := \|u\|_\infty + \|v\|_\infty$, where $\|w\|_\infty := \sup_{t \in [0, 1]} |w(t)|$.

In order to verify the existence of positive solutions for (S), we introduce now the following fixed-point theorem of cone expansion/compression type. We refer to [2, 3, 7] for proofs and further discussion of the fixed-point theory.

Lemma 2.1. *Let X be a Banach space with norm $|\cdot|$ and let $C \subset X$ be a cone in X . For $R > 0$, define $C_R = C \cap B[0, R]$, where $B[0, R] = \{x \in X : |x| \leq R\}$ denotes the closed ball of radius R centred at origin X . Assume that $F : C_R \rightarrow C$ is a compact map and that there exists $0 < r < R$ such that*

$$|Fx| \leq |x|, \quad x \in \partial C_r \quad \text{and} \quad |Fx| \geq |x|, \quad x \in \partial C_R,$$

or

$$|Fx| \geq |x|, \quad x \in \partial C_r \quad \text{and} \quad |Fx| \leq |x|, \quad x \in \partial C_R,$$

where $\partial C_R = \{x \in C : |x| = R\}$. Then F has a fixed point $u \in C$ with $r < |u| < R$.

Let us consider the cone C in X defined by

$$C = \{(u, v) \in X : (u, v)(0) = (u, v)(1) = 0 \text{ and } u, v \text{ are concave functions}\}.$$

Using the concavity of the functions $u(t)$ and $v(t)$, we may easily prove the following elementary result.

Lemma 2.2. *For each $(u, v) \in C$ and $(\alpha, \beta) \subset (0, 1)$, we have*

$$\inf_{t \in [\alpha, \beta]} (u(t) + v(t)) \geq \alpha(1 - \beta)\|(u, v)\|.$$

Lemma 2.3. *$F : X \rightarrow X$ is completely continuous and $F(C) \subset C$.*

Proof. We only give the main ideas of the proof. The Arzela–Ascoli theorem implies that A and B are concave functions and therefore $F : X \rightarrow X$ is completely continuous. It is easy to see that A and B (the coordinates functions of $F(u, v)$) are twice differentiable on $(0, 1)$ with $A'' \leq 0$ and $B'' \leq 0$. This implies that A and B are concave functions and therefore $F(C) \subset C$. □

3. Proof of Theorem 1.1

3.1. The existence of the first positive solution

Using the fact that f, g satisfy assumptions (H_0) , (H_1) and (H_3) , we apply Lemma 2.1 to prove the existence of a first positive solution for (S). Let $(a_1, b_1) \in [0, +\infty)^2 \setminus \{(0, 0)\}$ and $(a_2, b_2) \in (0, +\infty)^2$ be fixed.

Lemma 3.1. *Assume that conditions (H₀) and (H₁) hold. Then there exists R₁ > 0 such that, for all R ∈ (0, R₁], we have*

$$\|F(u, v)\| \geq \|(u, v)\| \quad \text{for each } (u, v) \in \partial C_R.$$

Proof. From (H₀), we have

$$\begin{aligned} f(t, u, v, a_1, b_1) &= d(t)h\left(\left(\frac{\eta_1}{\eta_2 - t}\right)^{1/(N-2)}, u + ta_1, v + tb_1\right) \\ &\geq d(t)h_1\left(\left(\frac{\eta_1}{\eta_2 - t}\right)^{1/(N-2)}, ta_1, tb_1\right). \end{aligned}$$

Thus, assuming (1.2), we obtain that there exist constants 0 < α₁ < β₁ < 1 such that

$$f_0 = \lim_{|(u,v)| \rightarrow 0} \frac{f(t, u, v, a_1, b_1)}{|(u, v)|} = +\infty \quad \text{uniformly for } t \in [\alpha_1, \beta_1]. \tag{3.1}$$

Now, using (3.1), for each M > 0, there exists R₁ > 0 such that

$$f(t, u, v, a_1, b_1) \geq M|(u, v)| \quad \text{for all } (u, v) \in [0, R_1]^2.$$

Therefore, for all (u, v) ∈ C_{R₁},

$$\begin{aligned} \|F(u, v)\| &\geq \|A(u, v)\|_\infty \\ &\geq a_2 \int_0^1 K\left(\frac{1}{2}, \tau\right) f(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau \\ &\geq a_2 M \int_{\alpha_1}^{\beta_1} K\left(\frac{1}{2}, \tau\right) [u(\tau) + v(\tau)] \, d\tau \\ &\geq a_2 \alpha_1 (1 - \beta_1) M \|(u, v)\| \int_{\alpha_1}^{\beta_1} K\left(\frac{1}{2}, \tau\right) \, d\tau. \end{aligned}$$

Finally, taking M > 0 sufficiently large, we conclude the proof of Lemma 3.1. □

Lemma 3.2. *Assume hypotheses (H₀) and (H₃). Then there exists R₂ > R₁ such that, for all R ≥ R₂, we have*

$$\|F(u, v)\| \leq \|(u, v)\| \quad \text{for each } (u, v) \in \partial C_R.$$

Proof. From assumptions (H₀) and (H₃), it is not difficult to see that

$$\lim_{|(u,v)| \rightarrow +\infty} \frac{f_1(r, u, v, a_1, b_1)}{|(u, v)|} = \lim_{|(u,v)| \rightarrow +\infty} \frac{g_1(r, u, v, a_1, b_1)}{|(u, v)|} = 0.$$

Thus, given δ > 0, there exists R₂ > R₁ such that, for all τ ∈ [0, 1] and |(u, v)| ≥ R₂,

$$f_1(\tau, u, v, a_1, b_1) \leq \delta |(u, v)| \quad \text{and} \quad g_1(\tau, u, v, a_1, b_1) \leq \delta |(u, v)|.$$

Thus, using (2.2), for all $(u, v) \in C_{R_2}$,

$$\begin{aligned} \|A(u, v)\|_\infty &= a_2 \int_0^1 K(t, \tau) f(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau \\ &\leq a_2 c_h \int_0^1 K(t, \tau) f_1(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau \\ &\leq a_2 c_h \int_0^1 K(t, \tau) f_1(\tau, \|u\|_\infty, \|v\|_\infty, a_1, b_1) \, d\tau \\ &\leq a_2 c_h \delta \|(u, v)\| \int_0^1 K(t, \tau) \, d\tau. \end{aligned}$$

Similarly, we may prove that

$$\|B(u, v)\|_\infty \leq a_2 c_h \delta \|(u, v)\| \int_0^1 K(t, \tau) \, d\tau.$$

Hence, taking $\delta > 0$ small enough, we have

$$\|F(u, v)\| = \|A(u, v)\|_\infty + \|B(u, v)\|_\infty \leq R_2 = \|(u, v)\|.$$

□

In view of Lemmas 3.1 and 3.2, as a direct consequence of Lemma 2.1, since $R_1 < R_2$, the proof of the first part of Theorem 1.1 is now complete.

3.2. The proof of the second part of Theorem 1.1

Lemma 3.3. *Assume that conditions (H_0) and (H_1) hold. Given $\bar{R} > 0$, there exists $\Lambda > 0$ such that, for all $(a_1, b_1) \in [0, +\infty)^2$ and $(a_2, b_2) \in (0, +\infty)^2$ with $\min\{a_2, b_2\} \geq \Lambda$, we have*

$$\|F(u, v)\| \geq \|(u, v)\| \quad \text{for each } (u, v) \in \partial C_{\bar{R}}, \tag{3.2}$$

where the constant Λ is independent of the parameters a_1 and b_1 .

Proof. We assume that (1.2) holds. Let $\alpha_1, \beta_1 \in (0, 1)$, as in the proof of Lemma 3.1, $\bar{R} > 0$ and $(u, v) \in C_{\bar{R}}$. According to assumption (H_0) and Lemma 2.2, we have

$$\begin{aligned} \|A(u, v)\|_\infty &\geq a_2 \int_{\alpha_1}^{\beta_1} K(\tfrac{1}{2}, \tau) f(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau \\ &\geq a_2 \int_{\alpha_1}^{\beta_1} K(\tfrac{1}{2}, \tau) f_1(\tau, u(\tau), v(\tau), a_1, b_1) \, d\tau \\ &\geq a_2 \int_{\alpha_1}^{\beta_1} K(\tfrac{1}{2}, \tau) f_1(\tau, \alpha_1(1 - \beta_1)\|u\|_\infty, \alpha_1(1 - \beta_1)\|v\|_\infty, 0, 0) \, d\tau. \end{aligned}$$

According to assumption (1.2), we see that

$$\bar{C} := \min\{f_1(\tau, \alpha_1(1 - \beta_1)u, \alpha_1(1 - \beta_1)v, 0, 0) : \tau \in [\alpha_1, \beta_1], u + v = \bar{R}\} > 0.$$

Thus we obtain that there exists a constant $\tilde{a} > 0$ such that

$$\|A(u, v)\|_\infty \geq a_2 \tilde{a}. \quad (3.3)$$

From (3.3), we get

$$\|F(u, v)\| \geq a_2 \tilde{a} \geq a_2 \frac{\tilde{a}}{\bar{R}} \bar{R}.$$

Hence there exists a positive constant Λ such that, for all $(a_1, b_1) \in [0, +\infty)^2$ and $(a_2, b_2) \in (0, +\infty)^2$ with $\min\{a_2, b_2\} \geq \Lambda$, we have

$$\|F(u, v)\| \geq \|(u, v)\| \quad \text{for each } (u, v) \in \partial C_{\bar{R}}. \quad (3.4)$$

We notice that Λ does not depend on the parameters a_1, b_1 . \square

Lemma 3.4. *Assume that (H_0) and (H_2) hold. Given $(a_2, b_2) \in (0, +\infty)^2$ with $\min\{a_2, b_2\} \geq \Lambda$, there exists $R_2 \in (0, \bar{R})$ and $\delta_1 > 0$ such that, for all $(a_1, b_1) \in [0, +\infty)^2$, with $a_1 + b_1 < \delta_1$, we have*

$$\|F(u, v)\| \leq \|(u, v)\| \quad \text{for each } (u, v) \in \partial C_{R_2}. \quad (3.5)$$

Proof. Using assumptions (H_0) – (H_2) , we have

$$\lim_{|(u, v, a_1, b_1)| \rightarrow 0} \frac{f(t, u, v, a_1, b_1)}{|(u, v, a_1, b_1)|} = 0.$$

Thus, given $\varepsilon > 0$, there exists $R_2 \in (0, \bar{R})$ such that, for all (u, v, a_1, b_1) with $u + v \leq R_2$ and $a_1 + b_1 \leq R_2$, we get

$$f(t, u, v, a_1, b_1) \leq \varepsilon |(u, v, a_1, b_1)|.$$

Let $(u, v) \in C_{R_2}$ and $a_1 + b_1 \leq R_2$. From the above estimate, we have

$$\begin{aligned} A(u, v)(t) &= a_2 \int_0^1 K(t, \tau) f(\tau, u(\tau), v(\tau)) \, d\tau \\ &\leq c_h a_2 \int_0^1 K(t, \tau) f_1(\tau, u(\tau), v(\tau)) \, d\tau \\ &\leq \varepsilon c_h a_2 (\|(u, v)\| + a_1 + b_1) \int_0^1 K(t, \tau) \, d\tau \\ &\leq \varepsilon c_h a_2 2R_2 \int_0^1 K(t, \tau) \, d\tau. \end{aligned}$$

Taking $\varepsilon > 0$ sufficiently small, we have

$$\|A(u, v)\|_\infty \leq \frac{1}{2} \|(u, v)\| \quad \text{for each } (u, v) \in \partial C_{R_2}.$$

We can now proceed analogously to prove that

$$\|B(u, v)\|_\infty \leq \frac{1}{2} \|(u, v)\| \quad \text{for each } (u, v) \in \partial C_{R_2}.$$

These two estimates together prove (3.5). \square

Finally, by considering $\bar{R} > R_2$ given in Lemmas 3.3 and 3.4, respectively, we apply once again Lemmas 3.1 and 3.2 to obtain R_3 and R_4 , with $0 < R_3 < R_2 < \bar{R} < R_4$, such that

$$\begin{aligned} \|F(u, v)\| &\geq \|(u, v)\|, & (u, v) \in \partial C_{R_3}, \\ \|F(u, v)\| &\leq \|(u, v)\|, & (u, v) \in \partial C_{R_2}, \\ \|F(u, v)\| &\geq \|(u, v)\|, & (u, v) \in \partial C_{\bar{R}}, \\ \|F(u, v)\| &\leq \|(u, v)\|, & (u, v) \in \partial C_{R_4}. \end{aligned}$$

Therefore, we can apply Lemma 2.1 to get three fixed points of F in C satisfying

$$R_3 < \|(u_1, v_1)\| < R_2 < \|(u_2, v_2)\| < \bar{R} < \|(u_3, v_3)\| < R_4,$$

and the proof is complete.

Acknowledgements. J.M.Ó. was supported by CNPq, PRONEX-MCT/Brazil and Millennium Institute for the Global Advancement of Brazilian Mathematics–IM-AGIMB. S.L. was supported by UTA grant 4734-02. P.U. was supported by DICYT–USACH.

Part of this work was done while the authors were visiting IMECC-UNICAMP/BRAZIL. The authors thank Professor Djairo de Figueiredo and all the faculty and staff of IMECC-UNICAMP for their kind hospitality.

References

1. C. BUNDLE AND L. A. PELETIER, Nonlinear elliptic problems with critical exponent in shrinking annuli, *Math. Ann.* **280** (1988), 1–19.
2. D. G. DE FIGUEIREDO, *Positive solutions of semilinear elliptic equations*, Lecture Notes in Mathematics, vol. 957, pp. 34–87 (Springer, 1982).
3. K. DEIMLING, *Nonlinear functional analysis* (Springer, 1985).
4. J. M. DO Ó, S. LORCA AND P. UBILLA, Local superlinearity for elliptic systems involving parameters, *J. Diff. Eqns*, in press.
5. D. R. DUNNINGER AND H. WANG, Existence and multiplicity of positive solutions for elliptic systems, *Nonlin. Analysis* **29** (1997), 1051–1060.
6. D. R. DUNNINGER AND H. WANG, Multiplicity of positive radial solutions for an elliptic system on an annulus, *Nonlin. Analysis* **42** (2000), 803–811.
7. D. GUO AND V. LAKSHMIKANTHAM, *Nonlinear problem in abstract cones* (Academic, 1988).
8. D. D. HAI, Positive solutions for semilinear elliptic equations in annular domains, *Nonlin. Analysis* **37** (1999), 1051–1058.
9. M. G. LEE AND S. S. LIN, On the positive solution for semilinear elliptic equations on annular domain with non-homogeneous Dirichlet boundary conditions, *J. Math. Analysis Applic.* **181** (1994), 348–361.
10. Y.-H. LEE, Multiplicity of positive radial solutions for multiparameter semilinear elliptic systems on an annulus, *J. Diff. Eqns* **174** (2001), 420–441.