

COMPOSITION IN FRACTIONAL SOBOLEV SPACES

HAIM BREZIS⁽¹⁾⁽²⁾ AND PETRU MIRONESCU⁽³⁾

1. Introduction

A classical result about composition in Sobolev spaces asserts that if $u \in W^{k,p}(\Omega) \cap L^\infty(\Omega)$ and $\Phi \in C^k(\mathbb{R})$, then $\Phi \circ u \in W^{k,p}(\Omega)$. Here Ω denotes a smooth bounded domain in \mathbb{R}^N , $k \geq 1$ is an integer and $1 \leq p < \infty$. This result was first proved in [13] with the help of the Gagliardo-Nirenberg inequality [14]. In particular if $u \in W^{k,p}(\Omega)$ with $kp > N$ and $\Phi \in C^k(\mathbb{R})$ then $\Phi \circ u \in W^{k,p}$ since $W^{k,p} \subset L^\infty$ by the Sobolev embedding theorem. When $kp = N$ the situation is more delicate since $W^{k,p}$ is not contained in L^∞ . However the following result still holds (see [2],[3])

Theorem 1. *Assume $u \in W^{k,p}(\Omega)$ where $k \geq 1$ is an integer, $1 \leq p < \infty$, and*

$$(1) \quad kp = N.$$

Let $\Phi \in C^k(\mathbb{R})$ with

$$(2) \quad D^j \Phi \in L^\infty(\mathbb{R}) \quad \forall j \leq k.$$

Then

$$\Phi \circ u \in W^{k,p}(\Omega)$$

The proof is based on the following

Lemma 1. *Assume $u \in W^{k,p}(\Omega) \cap W^{1,kp}(\Omega)$ where $k \geq 1$ is an integer and $1 \leq p < \infty$. Assume $\Phi \in C^k(\mathbb{R})$ satisfies (2). Then*

$$\Phi \circ u \in W^{k,p}(\Omega).$$

Acknowledgment: The first author (H.B.) is partially supported by a European Grant ERB FMRX CT98 0201. He is also a member of the Institut Universitaire de France. Part of this work was done when the second author (P.M.) was visiting Rutgers University; he thanks the Mathematics Department for its invitation and hospitality. We thank S. Klainerman for useful discussions.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\texttt{TeX}}$

Proof of Theorem 1. Since $u \in W^{k,p}$ we have

$$Du \in W^{k-1,p} \subset L^q$$

by the Sobolev embedding with

$$\frac{1}{q} = \frac{1}{p} - \frac{k-1}{N}.$$

Applying assumption (1) we find $q = N = kp$ and thus $u \in W^{1,kp}$. We deduce from Lemma 1 that $\Phi \circ u \in W^{k,p}$.

Proof of Lemma 1. Note that if $u \in W^{k,p} \cap L^\infty$ with $k \geq 1$ integer and $1 \leq p < \infty$ then $u \in W^{1,kp}$ by the Gagliardo - Nirenberg inequality [14]. Thus, Lemma 1 is a generalization of the standard result about composition. In fact, it is proved exactly in the same way as in the standard case (when $u \in W^{k,p} \cap L^\infty$). When $k = 2$ the conclusion is trivial.

Assume, for example that, $k = 3$, then

$$W^{3,p} \cap W^{1,3p} \subset W^{2,3p/2}$$

by the Gagliardo - Nirenberg inequality. Then

$$D^3(\Phi \circ u) = \Phi'(u)D^3u + 3\Phi''(u)D^2uDu + \Phi'''(u)(Du)^3,$$

and thus $\Phi \circ u \in W^{3,p}$ since

$$\begin{aligned} \int |D^2u|^p |Du|^p &\leq \left(\int |D^2u|^{3p/2} \right)^{2/3} \left(\int |Du|^{3p} \right)^{1/3} \\ &\leq C \|u\|_{W^{3,p}}^{p/2} \|u\|_{W^{1,3p}}^{3p/2}. \end{aligned}$$

A similar argument holds for any $k \geq 4$.

Starting in the mid-60's a number of authors considered composition in various classes of "Sobolev spaces" $W^{s,p}$, where $s > 0$ is a real number and $1 \leq p < \infty$. The most commonly used are the Bessel potential spaces $L^{s,p}(\mathbb{R}^N) = \{f = G_s * g; g \in L^p(\mathbb{R}^N)\}$ where $\widehat{G}_s = (1 + |\xi|^2)^{-s/2}$ and the Besov spaces $B_p^{s,p}(\mathbb{R}^N)$ (who's definition is recalled below when s is **not** an integer). It is well-known (see e.g. [1],[19] and [20]) that if k is an integer, $L^{k,p}$ coincides with the standard Sobolev space $W^{k,p}$; also if $p = 2$, the Bessel potential spaces $L^{s,2}$ and the Besov spaces $B_2^{s,2}$ coincide for every s non-integer and they are usually denoted by H^s . When $p \neq 2$ the spaces $L^{s,p}$ and $B_p^{s,p}$ are distinct.

The first result about composition in fractional Sobolev spaces seems to be due to Mizohata [12] for $H^s, s > N/2$. In 1970 Peetre [15] considered $B_p^{s,p} \cap L^\infty$ using interpolation

techniques; a very simple direct argument for the same class, $B_p^{s,p} \cap L^\infty$, was given by M. Escobedo [10] (see the proof of Lemma 2 below).

Starting in 1980 techniques of dyadic analysis and Littlewood-Paley decomposition à la Bony [5] were introduced. For example, Y. Meyer [11] considered composition in $L^{s,p}$ for $sp > N$; see also [16],[4],[9] for H^s with $s > N/2$ or for $H^s \cap L^\infty$, any $s > 0$. We refer to [17],[6],[7],[18] and their bibliographies for other directions of research concerning composition in Sobolev spaces.

In what follow we denote by $W^{s,p}(\Omega)$ the restriction of $B_p^{s,p}(\mathbb{R}^N)$ to Ω when s is not an integer. Our main result is the following

Theorem 2. *Assume $u \in W^{s,p}(\Omega)$ where $s > 1$ is a real number, $1 < p < \infty$, and*

$$(3) \quad sp = N.$$

Let $\Phi \in C^k(\mathbb{R})$, where $k = [s] + 1$, be such that

$$(4) \quad D^j \Phi \in L^\infty(\mathbb{R}) \quad \forall j \leq k.$$

Then

$$\Phi \circ u \in W^{s,p}(\Omega).$$

The proof of Theorem 2 relies on a variant of Lemma 1 for fractional Sobolev spaces.

Lemma 2. *Let $u \in W^{s,p}(\Omega)$, where $s > 1$ is a real number and $1 < p < \infty$. Assume, in addition, that $u \in W^{\sigma,q}$ for some $\sigma \in (0, 1)$ with*

$$(5) \quad q = sp/\sigma.$$

Let $\Phi \in C^k(\mathbb{R})$, where $k = [s] + 1$, be such that (4) holds. Then

$$\Phi \circ u \in W^{s,p}$$

Proof of Theorem 2. By the Sobolev embedding theorem we have

$$W^{s,p} \subset W^{r,q}$$

with $r < s$ and

$$\frac{1}{q} = \frac{1}{p} - \frac{(s-r)}{N}.$$

In view of assumption (3) we find

$$q = N/r.$$

In particular,

$$u \in W^{\sigma,q}$$

for **all** $\sigma \in (0, 1)$ with

$$q = \frac{N}{\sigma} = \frac{sp}{\sigma}.$$

Thus we may apply Lemma 2 and conclude that $\Phi \circ u \in W^{s,p}$.

Remark 1. Theorem 2 is known to be true when the Sobolev spaces $W^{s,p}$ are replaced by the Bessel potential spaces $L^{s,p}$ with $sp = N$; see D. Adams and M. Frazier [3]. Even though the two results are closely related it does not seem possible to deduce one from the other. Their argument relies on a variant of Lemma 2 for Bessel potential spaces:

Let $u \in L^{s,p} \cap L^{1,sp}$ where $s > 1$ is a real number and $1 < p < \infty$. Let Φ be as in Lemma 2. Then $\Phi \circ u \in L^{s,p}$.

Remark 2. The assumption in Lemma 2, $u \in W^{s,p} \cap W^{\sigma,q}$, with $q = sp/\sigma$ for some $\sigma \in (0, 1)$, is **weaker** than the assumption $u \in W^{s,p} \cap L^\infty$ but it is **stronger** than the assumption $u \in W^{1,sp}$; this is a consequence of Gagliardo - Nirenberg type inequalities (see e.g. the proof of Lemma D.1 in the Appendix D of [8]). It is therefore natural to raise the following:

Open Problem. Is the conclusion of Lemma 2 valid if one assumes only $u \in W^{s,p} \cap W^{1,sp}$ where $s > 1$ is a (non-integer) real number?

Before giving the proof of Lemma 2 we recall some properties of $W^{s,p}$ when s is not an integer.

When $0 < \sigma < 1$ and $1 < p < \infty$ the standard definition of $W^{\sigma,p}$ is

$$W^{\sigma,p}(\Omega) = \{f \in L^p(\Omega); \int \int \frac{|f(x) - f(y)|^p}{|x - y|^{N+\sigma p}} dx dy < \infty\}.$$

If $s > 1$ is not an integer write $s = [s] + \sigma$ where $[s]$ denotes the integer part of s and $0 < \sigma < 1$. Then

$$W^{s,p}(\Omega) = \{f \in W^{[s],p}(\Omega), D^\alpha f \in W^{\sigma,p} \text{ for } |\alpha| = [s]\}.$$

There is a very useful characterization of $W^{s,p}$ in terms of finite differences (see Triebel [20], p.110). Here it is more convenient to work with functions defined on all of \mathbb{R}^N and to consider their restrictions to Ω . Set

$$(\delta_h u)(x) = u(x + h) - u(x), \quad h \in \mathbb{R}^N,$$

so that

$$(\delta_h^2 u)(x) = u(x + 2h) - 2u(x + h) + u(x), \text{ etc...}$$

Given $s > 0$ not integer, fix **any** integer $M > s$. Then

$$W^{s,p} = \{f \in L^p; \int \int \frac{|\delta_h^M f(x)|^p}{|h|^{N+sp}} dx dh < \infty\}.$$

Proof of Lemma 2. It suffices to consider the case where s is not an integer. For simplicity we treat just the case where $1 < s < 2$. The same argument extends to general $s > 2$, s noninteger, using the same type of computations as in Escobedo [10].

The key observation is that $\delta_h^2(\Phi \circ u)$ can be expressed in terms of $\delta_h^2 u$ and $\delta_h u$. This is the purpose of our next computation.

Set

$$\begin{aligned} X &= u(x + 2h) \\ Y &= u(x + h) \\ Z &= u(x). \end{aligned}$$

Since $\Phi'' \in L^\infty(\mathbb{R})$ we have

$$(6) \quad \Phi(X) - \Phi(Y) = \Phi'(Y)(X - Y) + o(|X - Y|^2)$$

and since $\Phi' \in L^\infty(\mathbb{R})$ we also have

$$(7) \quad \Phi(X) - \Phi(Y) = \Phi'(Y)(X - Y) + o(|X - Y|).$$

Combining (6) and (7) we find

$$\Phi(X) - \Phi(Y) = \Phi'(Y)(X - Y) + o(|X - Y|^a)$$

for any $1 \leq a \leq 2$ (we will choose a specific value of a later) Similarly

$$\Phi(Z) - \Phi(Y) = \Phi'(Y)(Z - Y) + o(|Z - Y|^a)$$

Since

$$\delta_h^2(\Phi \circ u)(x) = (\Phi(X) - \Phi(Y)) + (\Phi(Z) - \Phi(Y)),$$

one finds

$$(8) \quad |\delta_h^2(\Phi \circ u)(x)| \leq C(|\delta_h^2 u(x)| + |\delta_h u(x + h)|^a + |\delta_h u(x)|^a).$$

This yields

$$(9) \quad \int \int \frac{|\delta_h^2(\Phi \circ u)(x)|^p}{|h|^{N+sp}} dx dh \leq C \int \int \frac{|\delta_h^2 u(x)|^p}{|h|^{N+sp}} dx dh + C \int \int \frac{|\delta_h u(x)|^{ap}}{|h|^{N+sp}} dx dh.$$

The first integral on the right-hand side of (9) is finite since $u \in W^{s,p}$. To handle the second integral we argue as follows. From the assumption $u \in W^{s,p} \cap W^{\sigma,q}$ with $\sigma \in (0,1)$ and q given by (5) we know that

$$(10) \quad \int \int \frac{|\delta_h^2 u(x)|^p}{|h|^{N+sp}} dx dh < \infty \text{ and } \int \int \frac{|\delta_h^2 u(x)|^q}{|h|^{N+sp}} dx dh < \infty.$$

From (10) and Hölder's inequality we derive that

$$(11) \quad \int \int \frac{|\delta_h^2 u(x)|^r}{|h|^{N+sp}} dx dh < \infty$$

for all $r \in [p, q]$, i.e., $u \in W^{\tau,r}$ with $\tau = sp/r$. We now choose

$$a = \min\{2, s/\sigma\}, \text{ so that } a \in [1, 2]$$

and $r = ap \in [p, q]$. It follows that

$$\int \int \frac{|\delta_h u(x)|^{ap}}{|h|^{N+sp}} dx dh < \infty,$$

which is the desired inequality.

Remark 3. There could be another natural proof of Theorem 2 by induction on $[s]$. One might attempt to prove that

$$D(\Phi \circ u) = \Phi'(u)Du \in W^{s-1,p}.$$

Note that $u \in W^{(s-1), N/(s-1)}$ and thus (by induction) we would have $\Phi'(u) \in W^{(s-1), N/(s-1)}$. On the other hand $Du \in W^{s-1,p}$. In order to conclude we need a lemma about products, but we are not aware of any such tool.

Remark 4. When s (or equivalently p) is a **rational** number, and $\Phi \in C^\infty$ with $D^j \Phi \in L^\infty \forall j$, there is a simple proof of Theorem 2 based on trace theory and Theorem 1. Assume for simplicity that $\Omega = \mathbb{R}^N$. Suppose that s is not an integer, but that $s_1 = s + 1/p$ is an integer. Then u is the trace of some function $u_1 \in W^{s_1,p}(\mathbb{R}^{N+1})$. Then $s_1 p = N+1$ and by Theorem 1 we deduce that $\Phi \circ u_1 \in W^{s_1,p}(\mathbb{R}^{N+1})$. Taking traces we find $\Phi \circ u \in W^{s,p}(\mathbb{R}^N)$. If s_1 is not an integer we keep extending u_1 to higher dimensions and stop at the first integer k such that $s_k = s + k/p$ is an integer (this is possible since p is rational and $s + k/p = (N+k)/p$ becomes an integer for some integer k). We have an extension $u_k \in W^{s_k,p}(\mathbb{R}^{N+k})$ of u . Then $\Phi \circ u_k \in W^{s_k,p}(\mathbb{R}^{N+k})$ by Theorem 1. Taking back traces yields $u \in W^{s,p}$.

REFERENCES

1. R. Adams, *Sobolev spaces*, Acad. Press, 1975.
2. D.R. Adams, *On the existence of capacitary strong type estimates in \mathbb{R}^n* , Ark. Mat. **14**, (1976), 125-140.
3. D.R. Adams and M. Frazier, *Composition operators on potential spaces*, Proc. Amer. Math. Soc., **114**, (1992), 155-165.
4. S. Alinhac and P. Gérard, *Opérateurs pseudo-différentiels et théorème de Nash-Moser*, Interéditions, 1991.
5. J.-M. Bony, *Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles nonlinéaires*, Ann. Sc. Ec. Norm Sup. **14**, (1981), 209-246.
6. G. Bourdaud, *Le calcul fonctionnel dans les espaces de Sobolev*, Invent. Math., **104**, (1991), 435-446.
7. G. Bourdaud and Y. Meyer, *Fonctions qui opèrent sur les espaces de Sobolev*, J. Funct. Anal., **97**, (1991), 351-360.
8. J. Bourgain, H. Brezis and P. Mironescu, *Lifting in Sobolev spaces*, J. d'Analyse **80**, (2000), 37-86.
9. J.-Y. Chemin, *Fluides parfaits incompressibles*, Astérisque **230**, 1995.
10. M. Escobedo, *Some remarks on the density of regular mappings in Sobolev classes of S^M -valued functions*, Rev. Mat. Univ. Complut. Madrid, **1**, (1988), 127-144.
11. Y. Meyer, *Remarques sur un théorème de J.-M. Bony*, Suppl. Rend. Circ. Mat. Palermo, Series II **1**, (1981), 1-20.
12. S. Mizohata, *Lectures on the Cauchy problem*, Tata Inst., Bombay, 1965.
13. J. Moser, *A rapidly convergent iteration method and non-linear differential equations*, Ann. Sc. Norm. Sup. Pisa **20**, (1966), 265-315.
14. L. Nirenberg, *On elliptic partial differential equations*, Ann. Sc. Norm. Sup. Pisa, **13**, (1959), 115-162.
15. J. Peetre, *Interpolation of Lipschitz operators and metric spaces*, Mathematica (Cluj) **12**, (1970), 1-20.
16. J. Rauch and M. Reed, *Nonlinear microlocal analysis of semilinear hyperbolic systems in one space dimension*, Duke Math J. **49**, (1982), 397-475.
17. T. Runst, *Mapping properties of non-linear operators in spaces of Triebel-Lizorkin and Besov type*, Analysis Mathematica, **12**, (1986), 313-346.
18. T. Runst and W. Sickel, *Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations*, Walter de Gruyter, Berlin and New York, 1996.
19. H. Triebel, *Interpolation theory, function spaces, differential operators*, North-Holland, Amsterdam, 1978.
20. H. Triebel, *Theory of function spaces*, Birkhäuser, Basel and Boston, 1983.

(1) ANALYSE NUMÉRIQUE
UNIVERSITÉ P. ET M. CURIE, B.C. 187
4 PL. JUSSIEU
75252 PARIS CEDEX 05

(2) RUTGERS UNIVERSITY
DEPT. OF MATH., HILL CENTER, BUSCH CAMPUS
110 FRELINGHUYSEN RD, PISCATAWAY, NJ 08854
E-mail address: brezis@ccr.jussieu.fr; brezis@math.rutgers.edu

(3) DEPARTEMENT DE MATHÉMATIQUES
UNIVERSITÉ PARIS-SUD
91405 ORSAY
E-mail address: : Petru.Mironescu@math.u-psud.fr